

Lack of Association of *CYP2E1* and *CYP1A1* Polymorphisms with Osteoporosis in Postmenopausal Women

Sadegh Fattahi¹, Gholam Ali Yousefi², Galia Amirbozorgi¹, Maryam Lotfi¹
Ali Naeiji¹, Mohsen Asouri¹, Saeid Kavosian¹, Effat Hayati¹, Ali Asghar Ahmadi¹
Haleh Akhavan-Niaki^{3*}

1- North Research Center, Pasteur Institute of Iran, Amol, Iran.

2- Radiology Department, Valiasr Hospital, Social Security Organization, GhaemShahr, Mazandaran, Iran.

3- Department of Genetics and Anatomy, Faculty of Medicine, Babol University of Medical Sciences, Babol, Iran.

Submitted 20 Oct 2015; Accepted 16 Nov 2015; Published 24 Dec 2015

Osteoporosis is a metabolic bone disease affecting mostly elderly women. As metabolizing enzymes, the roles of few cytochromes have been studied in osteoporosis development. The aim of this study was to assess for the first time the association of *CYP2E1* and *CYP1A1* polymorphisms and osteoporosis in postmenopausal women. 112 postmenopausal women presenting osteoporosis and 93 age and sex matched healthy controls originating from north Iran were enrolled in this study. Rs2031920 and rs3813867 at *CYP2E1* as well as rs4646421 and rs2198843 at *CYP1A1* loci were studied in all subjects using polymerase chain reaction and restriction fragment length polymorphism analysis. Genotype analysis for rs2031920 showed that the CT genotype was present only in osteoporotic patients with a frequency of 4.17%. Similarly GC genotype at rs3813867 locus was present only in osteoporotic patients with a frequency of 3.13%. [G; T] and [C; C] haplotypes for rs3813867- rs2031920 were found with low frequencies only in osteoporotic patients. TT genotype at rs4646421 locus was higher in osteoporotic (8.05%) versus control subjects (3.22%), 14.28% of cases were homozygous for the C allele at rs2198843 locus which is higher than controls (11.84%). The [C; G] haplotype for rs4646421- rs2198843 was predominant in cases (54.47%) and controls (58.6%). GT haplotype was rare in both groups. No significant differences in genotypes or haplotypes frequencies were observed among the osteoporotic and normal subjects. *CYP1A1* and *CYP2E1* loci are not associated to osteoporosis risk in the studied population.

Keywords: Osteoporosis; *CYP1A1*; *CYP2E1*; polymorphism

Osteoporosis is a silent metabolic bone disease characterized by low bone mass and reduction in bone mineral density (BMD) leading to increased susceptibility to fractures (1, 2). Different

factors such as environmental, genetic, gender, age, nutrition and hormones are associated with osteoporosis (3-6). Women are more affected than men and according to the World Health Organization

*Correspondence: Department of Genetics and Anatomy, Faculty of Medicine, Babol University of Medical Sciences, Babol, Iran.
E-mail: halehakhavan@yahoo.com

on, 25% of women and 12% of men become affected during their lives (7). Based on, 13% to 18% of women over 50 years have osteoporosis due to minimal ovarian activity that leads to decrease levels of estrogen in postmenopausal women and risk factor for bone loss. Genetic factors including Vitamin D receptor, Estrogen receptor, calcium-sensing receptor and cytochrome P450 genes have been reported as important players in the pathogenesis of osteoporosis (6, 8-10). Estrogen have essential role in bone turnover and establishing balance between bone formation and resorption. Moreover, metabolic pathways related to estrogen are important determinants of bone mineral density (BMD) in postmenopausal women. (11-14). The cytochrome P450 enzymes are a group of heme-containing enzymes placed in the endoplasmic reticulum of hepatocytes. They are required for the removal of carcinogenic compounds from the body (15). They are also essential in the metabolism of many medications, production of numerous agents such as cholesterol, steroids, vitamin D3 and vitamin K as well as metabolism and biosynthesis of estrogen. There are more than 50 CYP450 enzymes but six of them (CYP1A2, 2C9, 2C19, 2D6, 2E1 and 3A4) metabolize more than 85 percent of drugs and metabolites (16). Cyp1A1, Cyp19, Cyp17, CYP3A4, CYP1A2 and CYP1B1 have been demonstrated to have a role in osteoporosis development (9, 17, 18). Cytochrome P450 2E1 (CYP2E1), a member of the cytochrome P450 superfamily is important for the metabolic activation of many low-molecular-weight toxicants such as N-nitrosamines, aniline, vinyl chloride, urethane and alcohol that is expressed primarily in the liver (19). It has been reported that *CYP2E1* polymorphism is involved in several types of diseases including cancer, hepatitis and tuberculosis (20). Recently, it has been reported that RsaI/PstI polymorphism as well as sex hormones may influence human *CYP2E1*

gene expression (21). Also, it has been reported that *CYP1A1* polymorphism may play an important role in estrogen metabolism (22-24). The aim of this study was to investigate for the first time the association between *CYP2E1* and *CYP1A1* polymorphisms with osteoporosis in a population of postmenopausal women originating from north Iran.

Materials & Methods

Subjects

This case-control study contained 112 postmenopausal women presenting osteoporosis and 93 age and sex matched healthy controls that were referred to Valiasr Hospital in Ghaemshar, Mazandaran, Iran. All subjects consented to participate to this study and demographic data were collected upon personal interview. Also, the study was approved by the scientific and ethics committees of north research center of Pasteur institute, Amol, Iran.

DNA extraction

Genomic DNA was extracted from whole blood using salting out method. Briefly, 5ml whole blood was digested with proteinase K followed by 6 M sodium acetate treatment. The DNA was collected after alcohol precipitation, and then dissolved in 10 mM Tris (pH 8.0) and 0.1 mM EDTA.

Analysis of *CYP2E1* gene polymorphisms

Two single-nucleotide polymorphisms of *CYP2E1* (rs3813867 and rs2031920) were studied by polymerase chain reaction-restriction fragment length polymorphism (PCR-RFLP) analysis. The cycling conditions consisted of an initial denaturation at 94 °C for 5 min followed by 35 cycles, each consisting of denaturation at 94 °C for 45 seconds, annealing at 59°C for 1 min, extension at 72 °C for 60 seconds and one final cycle of extension at 72°C for 7 min. The PCR products were

Table 1. PCR and RFLP conditions for the *CYP11A1* and *CYP2E1* genotyping

Gene	SNP	Primer sequence	PCR product (bp)	Enzyme	RFLP product (bp)	Reference or source
<i>CYP11A1</i>	rs4646421	F: 5'-GCTCAATCTAGCTGG TCTCCA -3' R: 5'-GAGTCCTACCCACCA CACTTAG -3'	417	NspI	231, 138	This study
<i>CYP11A1</i>	rs2198843	F: 5'-GTCCCTCAGAGAACA AGGCAG -3' R: 5'- CTCCTCTCAGCTTC ACAGGCA -3'	404	PstI	299, 105	This study
<i>CYP2E1</i>	rs2031920	F: 5'-CCA GTC GAG TCT ACA TTG TCA-3'	413	RsaI	354, 59	(25)
	rs3813867	R: 5'-TTC ATT CTG TCT TCT AAC TGG-3'	413	PstI	299, 114	

digested overnight with either RsaI (rs2031920) or PstI (rs3813867) restriction enzymes at 37°C. The evaluation of the digestion patterns was achieved by electrophoresis on a 3% agarose gel followed by ethidium bromide staining. The expected sizes of the products after digestion with RsaI were as follows: 413 bp for the T allele (no cutting), 354 bp and 59 bp for the C allele. The expected sizes after digestion with PstI are 299 bp and 114 bp for the C allele, while the G allele remains uncut with 413 bp size.

Analysis of *CYP11A1* gene polymorphisms

We developed specific PCR–restriction fragment length polymorphism tests for two single-nucleotide polymorphisms of *CYP11A1* (rs4646421 and rs2198843). PCRs for both SNPs were performed under following conditions, after initial denaturation at 95 °C for 5 minutes 35 cycles each consisting of denaturation at 94 °C for 45 seconds, annealing at 60 °C for 45 seconds, extension at 72 °C for 45 seconds and a final elongation step at 72 °C for 7 minutes were performed. The PCR products were digested overnight with either NspI (rs4646421) or PstI (rs2198843) restriction enzyme at 37°C and the presence of restriction sites was identified by agarose gel electrophoresis. The information about sequence of primer pairs, size of PCR products and digested products with restriction enzymes is listed in Table 1.

Statistical analyzes

The χ^2 test was used to evaluate differences in the distributions of genotypes and allele frequencies between cases and controls. Odds ratios (ORs) and 95% confidence intervals (95% CIs) were used to describe the strength of associations. An adjusted two-tailed P value less than 0.05 at 95% CI was considered as statistically significant.

Results

Age and BMD analysis

In this study, we examined 112 postmenopausal cases with osteoporosis aged 59.35 ± 7.76 years and 93 normal subjects aged 57.37 ± 5.56 years. The BMD analyzes showed that BMD was significantly lower in osteoporosis patients (0.817 ± 0.074) than normal subjects (1.231 ± 0.104) ($P < 0.0001$).

CYP2E1 genotyping

The frequencies of C and T alleles for rs2031920 as well as genotype distribution among studied subjects are shown in Table 2. The C allele was present in all normal subjects while its frequency was 0.982 among osteoporotic subjects. The wild type homozygous genotype (CC) was the most frequent in osteoporotic subjects (96.42%) as well as in normal subjects (100%). None of the analyzed subjects showed TT genotype. RsaI polymorphism genotype distributions were in Hardy-Weinberg equilibrium ($P = 0.847$). Figure 1A

Table 2. Distribution of CYP2E1 and CYP1A1 genotype and allele frequencies in osteoporotic cases compared to controls for rs2031920, rs3813867, rs4646421 and rs2198843

			wild type homozygous	Heterozygous	Mutant homozygous	OR (95% CI)
rs2031920 polymorphism	C allele	T allele	CC	CT	TT	
Osteoporosis (n=112)	0.982	0.018	108(96.42)	4(3.58)	0	1.00
Normal (n=93)	1	0	93(100)	0	0	7.75(0.41-145)
rs3813867 polymorphism	G allele	C allele	GG	GC	CC	
Osteoporosis (n=112)	0.986	0.014	109(97.32)	3(2.68)	0	1.00
Normal (n=93)	1	0	93(100)	0	0	5.9772 (0.30-117)
rs4646421 polymorphism	C allele	T allele	CC	CT	TT	1.00
Osteoporosis (n=112)	0.767	0.233	69(61.6%)	34(30.35 %)	9(8.05%)	1.1498(0.62-2.11)
Normal (n=93)	0.822	0.178	63(67.6%)	27(29.09%)	3(3.22%)	2.7391(0.70-10.57)
rs2198843 polymorphism	G allele	C allele	GG	GC	CC	1.00
Osteoporosis (n=112)	0.674	0.326	55(49.10 %)	41(36.62%)	16(14.28 %)	1.1069(0.60-2.01)
Normal (n=93)	0.704	0.296	49(52.68 %)	33(35.48 %)	11(11.84%)	1.2959(0.54-3.05)

represents the electrophoresis pattern after digestion of PCR products with RsaI.

The distributions of the genotypes and frequencies of C and G alleles for rs3813867 of the studied subjects are summarized in Table 2. The allele frequency for the G allele among osteoporotic population was 0.986, while all control subjects had the G allele. The wild type homozygous genotype (GG) was the most frequent in both groups. None of the analyzed subjects showed CC genotype. PstI polymorphism genotype distributions were in Hardy-Weinberg equilibrium (P= 0.885). Figure 1A is a representative of PstI genotyping analysis on PCR products of CYP2E1 gene.

CYP1A1 genotyping

The frequencies of the C and T alleles and genotype distribution of rs4646421 are summarized in Table 2. The C and T allele frequencies were found to be 0.822 and 0.178 among the control subjects, and 0.767 and 0.233 among osteoporotic subjects, respectively. The distribution of CC and CT genotypes in osteoporotic cases were 61.6% and 30.35%, respectively compared to 67.6% and 29.09% in healthy controls. TT genotype was higher in osteoporotic subjects (8.05%) versus control subjects (3.22%), but the difference was not significant. Observed genotypes were in Hardy-Weinberg equilibrium (P= 0.1401). Figure 1B shows




Figure 1. Representative electrophoresis patterns for *CYP3E1* and *CYP1A1* genotyping. A, RFLPs of the PCR products of *CYP2E1* gene digested with *Rsa* I and *Pst* I. Lane 1: Molecular Weight Marker (100 bp ladder); lanes 2 and 8: undigested; lanes 3 and 6 heterozygous; lanes 4 and 7: homozygous wild-type; lane 5: mutant homozygous. B, RFLPs of the PCR products of *CYP1A1* rs4646421 digested with *Nsp*I. Lane 1: Molecular Weight Marker (100 bp ladder); lane 2: mutant homozygous; lane 3: homozygous wild-type; lane 4: heterozygous. C, RFLPs of the PCR products of *CYP1A1* rs2198843 digested with *Pst*I. Lane 1: Molecular Weight Marker (100 bp ladder); lane 2: mutant homozygous; lanes 3 and 4: homozygous wild-type; lane 5: heterozygous.

-ws the electrophoresis pattern after digestion of PCR products with *Nsp*I.

The frequencies of the C and G alleles as well as genotype distribution in controls and patients for rs2198843 are shown in Table 2. Allelic frequencies for G and C allele were 0.674 and 0.326 among osteoporotic subjects versus 0.704 and 0.296 in control subjects, respectively. The distribution of GG and GC genotypes in osteoporotic cases was 49.1% and 36.62% compared to 52.68% and 35.48% in healthy controls, respectively. As shown in Table 1, 14.28% of osteoporotic cases were homozygous for the C allele which is higher than controls (11.84%) but the difference is not significant. Genotype distributions of rs2198843 was in Hardy-Weinberg equilibrium ($P=0.1401$). The electrophoresis pattern after digestion of PCR products with *st*I is shown in figure 1C.

Haplotypes analyzes

Haplotype data for rs2031920 and rs3813867 at *CYP2E1* locus are provided in Table 3. Our study revealed that the osteoporotic population has three haplotypes, while healthy controls have only one haplotype. But no significant difference in distribution of the common haplotype was observed in patients with osteoporosis and healthy controls. The predominant haplotype in this study was [G; C] Table 3.

Haplotype data for rs4646421 and rs2198843 at *CYP1A1* locus are shown in Table 4. Four haplotypes were identified in osteoporotic as well as normal subjects. The haplotype [C; G] was predominant in osteoporotic (54.47%) and control subjects (58.6%). In this study, [G; T] haplotype was rare in both osteoporotic and normal subjects.

Table 3 Associations between [rs3813867; rs2031920] haplotypes of *CYP2E1* and osteoporosis

	Osteoporosis n (%)*	Normal n (%)	OR (95% CI)	P value
[G; C]	218 (97.32)	186 (100)	1.00 (reference)	
[G; T]	1(0.446)	0	2.5606(0.10-63.23)	0.56
[C; C]	1(0.446)	0	2.5606(0.10-63.23)	0.56

*For 1.8% of the analyzed subjects haplotype assignment was not possible due to heterozygosity in both loci.

Table 4. Associations between [rs3813867; rs2031920] haplotypes of *CYP1A1* and osteoporosis

	Osteoporosis no. (%) [*]	Normal no. (%) [*]	OR (95% CI)	P value
[C; G]	122(54.47%)	109(58.6%)	1.00	
[C; C]	26(11.6%)	24(12.9%)	1.0332(0.56-1.90)	0.91
[T; G]	5(2.24%)	3(1.62%)	1.4891(0.34-6.37)	0.59
[T; C]	23(10.27%)	10(5.38%)	2.0549(0.93-4.51)	0.07

^{*}For 21.42 % of the analyzed osteoporotic subjects and 21.5% of normal subjects haplotype assignment was not possible due to heterozygosity in both loci.

Discussion

Many genetic factors were reported as having a significant effect on bone remodeling and bone mass. Therefore, polymorphism in genes controlling drugs and metabolism such as *CYP450* may act as strong candidate loci for increased risk of genetically influenced disorders such as osteoporosis. Several studies show that the genetic polymorphisms in *CYP450* are associated with osteoporosis in postmenopausal women (9, 17, 18, 26, 27). However, there is no information about the impact of *CYP2E1* (RsaI/ PstI) genetic polymorphisms on osteoporosis susceptibility. This is the first study evaluating the *CYP2E1* polymorphisms in osteoporotic subjects. In this study, rs2031920 and rs3813867 were studied in 93 healthy subjects and 112 patients with osteoporosis. Genotype analysis for rs2031920 showed that the CT genotype was present only in osteoporotic patients with a frequency of 3.58%. Genotype analysis for rs3813867 showed that the GC genotype was present only in osteoporosis patients with a frequency of 2.68%. There were no significant differences in genotype frequencies among the osteoporotic and normal subjects although the risk for the heterozygous genotype was seven-fold greater than the homozygous wild type CC genotype for rs2031920 (OR= 7.75) in osteoporotic cases. Similarly, the risk for the heterozygous genotype was five-fold greater than the homozygous wild type

GG genotype for rs3813867 (OR= 5.9772) in osteoporotic cases. Haplotypes distribution at *CYP2E1* locus was not significantly different between the osteoporotic subjects and controls (P= 0.5), but the risk for the [G; T] and [C; C] haplotype subjects to have osteoporosis is three-fold greater than it is for the [G; C] haplotype (OR= 2.56). Our findings on alleles and genotype frequencies are similar to another study in Iran which analyzed two hundred healthy individuals from the southwest population (28). The comparison of genotype distribution between Iranian and other populations shows that its distribution is similar to Indian, Turkish and some European populations such as German, British and French (21, 29-32) but is different from other Asians, including Japanese and Chinese and also from Italians (33-35). Based on WHO criteria, 11.9% of Native American women, 10% of Asians, 9.8% of Hispanics, 7.2% of Whites, and 4.2% of Blacks were osteoporotic (36). These distribution differences among ethnicities strongly support the genetic contribution to disease occurrence. Cyp17 and CYP 19 enzymes are involved in estrogen biosynthesis while CYP3A4, CYP1A2, CYP1B1 and CYP1A1 contribute to its metabolism (37). Polymorphism in the cytochrome P450 enzymes related to estrogen biosynthesis and metabolism may lead to increased risk of hormone-dependent disorders such as breast cancer (23, 24, 38-41) and osteoporosis (9, 17, 18, 26). It has been reported that *CYP1A1* polymorphism may play an

important role in estrogen metabolism (9, 22, 24, 42). Genotype analysis for rs4646421 showed that the frequency of TT genotype was higher in patients with osteoporosis than in controls. Although no significant associations were observed between osteoporotic and normal subjects, but osteoporosis risk was three times greater in individuals with TT genotype than individuals with CC and CT genotypes. Genotype analysis for rs2198843 showed that the distribution of genotypes was approximately similar in both groups. There were no significant differences in genotype frequencies among the osteoporotic subjects and normal ones. Haplotypes were neither significantly different between them ($P=0.07$), but the risk for the mutant [T; C] haplotype subjects to have osteoporosis was two times greater than it was for the wild haplotype [C; G] (OR= 2.05). In this study, we found [G; T] and [C; C] haplotypes for *CYP2E1* [rs3813867; rs2031920] and [C; T] haplotypes for *CYP1A1* [rs4646421; rs2198843] with low frequencies in osteoporotic patients. However, as osteoporosis is a complex disease and many genetic factors may contribute in the development of this disorder, a single polymorphism may not show a significant association with the disease unless it is strongly associated. Thus, other polymorphisms in candidate genes should be studied in order to find a polymorphism panel for osteoporosis risk evaluation. Correspondingly, it was also recommended to study a larger sample size, as minor alleles with strong impact on disease development may have more chances to be identified. Moreover, other population studies should be performed to replicate our findings.

Acknowledgements

We would like to express our gratitude to all patients who accepted to participate to this study. Authors are also grateful to Valiasr Hospital staff for

their assistance.

Conflict of interest

The authors declared no conflict of interests.

References

1. Stewart T L, Ralston S H. Role of genetic factors in the pathogenesis of osteoporosis. *J Endocrinol.* 2000 Aug;166(2):235-45.
2. Rizzoli R, Bonjour J P, Ferrari S L. Osteoporosis, genetics and hormones. *J Mol Endocrinol.* 2001 Apr;26(2):79-94.
3. Hamidi Z, Majdzadeh R, Soltani A A F, et al. Casual decomposition of risk factors in osteoporosis burden. *J Med Counc IRI.* 2007;24(4):381-92.
4. Ozbas H, Tutgun Onrat S, Ozdamar K. Genetic and environmental factors in human osteoporosis. *Mol Biol Rep.* 2012 Dec;39(12):11289-96.
5. Shao H, Tao M, Fan Y, et al. Vitamin d levels and other factors related to bone mineral density during pregnancy. *Aust N Z J Obstet Gynaecol.* 2012 Dec;52(6):571-5.
6. Zajickova K, Zofkova I. Osteoporosis: Genetic analysis of multifactorial disease. *Endocr Regul.* 2003 Mar;37(1):31-44.
7. Khan A. Advances in osteoporosis therapy. 2003 update of practical guidelines. *Can Fam Physician.* 2003 Apr;49:441-7.
8. Lambertini E, Penolazzi L, Sollazzo V, et al. Modulation of gene expression in human osteoblasts by targeting a distal promoter region of human estrogen receptor-alpha gene. *J Endocrinol.* 2002 Mar;172(3):683-93.
9. Napoli N, Villareal D T, Mumm S, et al. Effect of *cypl1a1* gene polymorphisms on estrogen metabolism and bone density. *J Bone Miner Res.* 2005 Feb;20(2):232-9.
10. Fang Y, van Meurs J B, d'Alesio A, et al. Promoter and 3'-untranslated-region haplotypes in the vitamin d receptor gene predispose to osteoporotic fracture: The rotterdam study. *Am J*

Hum Genet. 2005 Nov;77(5):807-23.

11. Abdallah B M, Bay-Jensen A C, Srinivasan B, et al. Estrogen inhibits dlk1/fa1 production: A potential mechanism for estrogen effects on bone turnover. *J Bone Miner Res.* 2011 Oct;26(10):2548-51.

12. Confavreux C B, Fontana A, Guastalla J P, et al. Estrogen-dependent increase in bone turnover and bone loss in postmenopausal women with breast cancer treated with anastrozole. Prevention with bisphosphonates. *Bone.* 2007 Sep;41(3):346-52.

13. Lin S Q. Effects of estrogen on bone-turnover and its prevention and control of osteoporosis. *Zhonghua Yi Xue Za Zhi.* 2005 Mar 23;85(11):728-9.

14. Napoli N, Armamento-Villareal R. Estrogen hydroxylation in osteoporosis. *Adv Clin Chem.* 2007;43:211-27.

15. Guengerich F P. Characterization of human cytochrome p450 enzymes. *FASEB J.* 1992 Jan 6;6(2):745-8.

16. Shapiro L E, Shear N H. Drug interactions: Proteins, pumps, and p-450s. *J Am Acad Dermatol.* 2002 Oct;47(4):467-84; quiz 85-8.

17. Somner J, McLellan S, Cheung J, et al. Polymorphisms in the p450 c17 (17-hydroxylase /17,20-lyase) and p450 c19 (aromatase) genes: Association with serum sex steroid concentrations and bone mineral density in postmenopausal women. *J Clin Endocrinol Metab.* 2004 Jan;89(1):344-51.

18. Koudu Y, Onouchi T, Hosoi T, et al. Association of cyp19 gene polymorphism with vertebral fractures in japanese postmenopausal women. *Biochem Genet.* 2012 Jun;50(5-6):389-96.

19. Lai C, Shields P G. The role of interindividual variation in human carcinogenesis. *J Nutr.* 1999 Feb;129(2S Suppl):552S-55S.

20. Roy B, Ghosh S K, Sutradhar D, et al. Predisposition of antituberculosis drug induced

hepatotoxicity by cytochrome p450 2e1 genotype and haplotype in pediatric patients. *J Gastroenterol Hepatol.* 2006 Apr;21(4):784-6.

21. Konstandi M, Cheng J, Gonzalez F J. Sex steroid hormones regulate constitutive expression of cyp2e1 in female mouse liver. *Am J Physiol Endocrinol Metab.* 2013 May 15;304(10):E1118-28.

22. Badawi A F, Cavalieri E L, Rogan E G. Role of human cytochrome p450 1a1, 1a2, 1b1, and 3a4 in the 2-, 4-, and 16alpha-hydroxylation of 17beta-estradiol. *Metabolism.* 2001 Sep;50(9):1001-3.

23. Hong C C, Thompson H J, Jiang C, et al. Association between the t27c polymorphism in the cytochrome p450 c17alpha (cyp17) gene and risk factors for breast cancer. *Breast Cancer Res Treat.* 2004 Dec;88(3):217-30.

24. Kisselev P, Schunck W H, Roots I, et al. Association of cyp1a1 polymorphisms with differential metabolic activation of 17beta-estradiol and estrone. *Cancer Res.* 2005 Apr 1;65(7):2972-8.

25. Hayashi S, Watanabe J, Kawajiri K. Genetic polymorphisms in the 5'-flanking region change transcriptional regulation of the human cytochrome p450iie1 gene. *J Biochem.* 1991 Oct;110(4):559-65.

26. Kang Y S, Park S Y, Yim C H, et al. The cyp3a4*18 genotype in the cytochrome p450 3a4 gene, a rapid metabolizer of sex steroids, is associated with low bone mineral density. *Clin Pharmacol Ther.* 2009 Mar;85(3):312-8.

27. Masi L, Becherini L, Gennari L, et al. Polymorphism of the aromatase gene in postmenopausal italian women: Distribution and correlation with bone mass and fracture risk. *J Clin Endocrinol Metab.* 2001 May;86(5):2263-9.

28. Shahriary G M, Galehdari H, Jalali A, et al. Cyp2e1*5b, cyp2e1*6, cyp2e1*7b, cyp2e1*2, and cyp2e1*3 allele frequencies in iranian populations. *Asian Pac J Cancer Prev.* 2012;13(12):6505-10.

29. Yang B, O'Reilly D A, Demaine A G, et al. Study of polymorphisms in the cyp2e1 gene in

- patients with alcoholic pancreatitis. *Alcohol*. 2001 Feb;23(2):91-7.
30. Neuhaus T, Ko Y D, Lorenzen K, et al. Association of cytochrome p450 2e1 polymorphisms and head and neck squamous cell cancer. *Toxicol Lett*. 2004 Jun 15;151(1):273-82.
31. Bouchardy C, Hirvonen A, Coutelle C, et al. Role of alcohol dehydrogenase 3 and cytochrome p-4502e1 genotypes in susceptibility to cancers of the upper aerodigestive tract. *Int J Cancer*. 2000 Sep 1;87(5):734-40.
32. Ulusoy G, Arinc E, Adali O. Genotype and allele frequencies of polymorphic cyp2e1 in the turkish population. *Arch Toxicol*. 2007 Oct;81(10):711-8.
33. Boccia S, Cadoni G, Sayed-Tabatabaei F A, et al. Cyp1a1, cyp2e1, gstm1, gstt1, ephx1 exons 3 and 4, and nat2 polymorphisms, smoking, consumption of alcohol and fruit and vegetables and risk of head and neck cancer. *J Cancer Res Clin Oncol*. 2008 Jan;134(1):93-100.
34. Quinones L, Lucas D, Godoy J, et al. Cyp1a1, cyp2e1 and gstm1 genetic polymorphisms. The effect of single and combined genotypes on lung cancer susceptibility in chilean people. *Cancer Lett*. 2001 Dec 10;174(1):35-44.
35. Sangrajrang S, Jedpiyawongse A, Srivatanakul P. Genetic polymorphisms of cyp2e1 and gstm1 in a thai population. *Asian Pac J Cancer Prev*. 2006 Jul-Sep;7(3):415-9.
36. Handa R, Ali Kalla A, Maalouf G. Osteoporosis in developing countries. *Best Pract Res Clin Rheumatol*. 2008 Aug;22(4):693-708.
37. Napoli N, Armamento-Villareal R. Effects of polymorphisms of the cyp450 enzyme genes on estrogen status and the risk for osteoporosis. *Curr Pharmacogenomics*. 2007;5(1):1-10.
38. De Vivo I, Hankinson S E, Li L, et al. Association of cyp1b1 polymorphisms and breast cancer risk. *Cancer Epidemiol Biomarkers Prev*. 2002 May;11(5):489-92.
39. Kristensen V N, Harada N, Yoshimura N, et al. Genetic variants of cyp19 (aromatase) and breast cancer risk. *Oncogene*. 2000 Mar 2;19(10):1329-33.
40. Taioli E, Bradlow H L, Garbers S V, et al. Role of estradiol metabolism and cyp1a1 polymorphisms in breast cancer risk. *Cancer Detect Prev*. 1999;23(3):232-7.
41. Watanabe J, Shimada T, Gillam E M, et al. Association of cyp1b1 genetic polymorphism with incidence to breast and lung cancer. *Pharmacogenetics*. 2000 Feb;10(1):25-33.
42. Sowers M R, Wilson A L, Kardia S R, et al. Cyp1a1 and cyp1b1 polymorphisms and their association with estradiol and estrogen metabolites in women who are premenopausal and perimenopausal. *Am J Med*. 2006;119(9):S44-S51.