Investigation on Effects of Parenterally Given Vitamin B Complex on Ruminal Protozoa in Cattle

Bulent Elitok¹*, Seda Akgun²

- 1. Afyon Kocatepe University, Faculty of Veterinary Medicine, Department of Internal Medicine, Afyonkarahisar, Turkey.
- 2. Provincial Directorate of Agriculture, Afyonkarahisar, Turkey.

Submitted 15 Aug 2017; Accepted 27 Sep 2017; Published 29 Oct 2017

B complex vitamins play an important role in the metabolism of carbohydrates, proteins and fats. The aim of this study was to determine the effects of B complex vitamins administered by exogenous route on rumen protozoa, hematological and blood biochemical parameters in clinically healthy cattles. Thirty cattles aged between 6 to 18 months, and breeding in Afyonkarahisar Province and surrounding regions were studied. Ten clinically healthy animals served as control group. Twenty study group animals were given 10-20 ml B complex vitamins, 3 times during 3 days, by intramuscular route. Clinical (body temperature, pulse and respiration rates, rumen contractions), hematological (total leukocyte count, erythrocyte count, hemoglobin and hematocrit measurements), and serum biochemical (aspartate aminotransferase, glucose, total protein, albumin) parameters along with rumen protozoa status were measured in all animals. Regarding hematological parameters, there was no significant difference between the study groups in term of time intervals (P> 0.05). The mean values of WBC, RBC, neutrophils, HGB, HTC, MCV, MCH and MCHC were different between the case and the control groups (P< 0.05), but the mean values of lymphocytes, monocytes, and basophils were not different (P> 0.05). AST, SDH, ALP enzyme levels averaged within the reference limits, but were significantly higher in the case group (P< 0.05). Consequently, the use of B complex vitamins have proved to be of great benefit as it did not caused adverse effects on the digestive system at certain intervals in cattle.

Keywords: Afyonkarahisar, B complex vitamins, cattle, rumen, protozoa

Rumen produced B complex vitamins play an important role in the metabolism of carbohydrates, proteins and fats (1-3). Ruminants remove their needs by digesting bacteria found in rumen or by taking B vitamins that are free in the body fluid (4-7). B complex vitamins include different vitamins such as thiamin (vitamin B1), riboflavin (vitamin B2), niacin (vitamin B3 or PP), choline (vitamin B4), pantothenic acid (vitamin B5), pyridoxine (vitamin B6), biotin, cyanocobalamine (vitamin B12). Vitamin B1 plays a key role in energy metabolism, and is destroyed by an enzyme called

thiaminase, which is secreted by some bacteria in the rumen (8). When the thiamin level falls, neuro-transmitters do not form, and communication between the nerves may be interrupted (9-12). The lack of vitamin B1 is very common in ruminants as oral antibiotics, sulfa group medications, and antacids can cause thiamine deficiency (1, 13). Vitamin B2 is crucial for some enzymes for fatty acid and amino acid synthesis. In the absence of riboflavin, non-specific symptoms such as anorexia, chronic diarrhea, and growth regression are seen (13). Niacin plays a critical role in oxidation and

glucose synthesis of rumen-forming volatile fatty acids (VFAs), and inhibits the formation of ketosis (14). Choline is the antioxidant of acetylcholine, which indirectly plays an effective role in fat metabolism by increasing carnitine synthesis (15). Vitamin B5 is an obligatory vitamin in coenzyme A production, and plays a vital role as a catalyst in the conversion of carbohydrates, fats and proteins into energy (4, 16-19). Pyridoxine is absorbed from the ileum by passive diffusion when it is produced in rumen. Absence of this vitamin causes axonal degeneration, demyelination in the peripheral nerves, cramps, inadequate growth, and anemia (20, 21). In cattle, dietary biotin (vitamin B8) supplement has been shown to improve nail wall damages and many nail diseases caused by wet soils, eliminate pregnancy and lameness problems, and improve liver fatigue (7, 21). In absence of folic acid (vitamin B9), DNA synthesis is disturbed, and conversion of glycine and serine to each other becomes more difficult (5, 7, 18). In ruminants, the absorption of vitamin B12 has many beneficial effects such as destruction of an important amount of dietary cobalt by the rumen microflora (7), food proteins digestion and vitamin B12 release for the production of internal factors by a functional abomasum, and trypsin secretion by a functional pancreas for protein digestion (13, 22).

Hematological and blood biochemical parameters as well as rumen fluid analyzes provide important information about the animal's health status (23-25). To the best of our knowledge, there is no literature report directly investigating the effect of B complex vitamins administered by the parenteral route on the content of rumen in cattle, although there are some studies examining the effects of orally administered B complex vitamins on the rumen protozoa (11, 12).

In this study, we aimed to determine the effects of B complex vitamins administered by intramuscular (IM) route on rumen protozoa, hematological and blood biochemical parameters, in clinically healthy cattles.

Materials and methods

Study design

The study was carried out in Afyonkarahisar Province on 30 cattles (6 to 18 months old) who were breeding public (study group). Ten clinically healthy animals having the same age range and nutritional requirements were selected as control group. 10-20 ml B complex vitamins (MUL-TİKOM-B, Celikler Pharmaceuticals and Trade Co., Ankara / Turkey) were given by IM route for 3 days to twenty case group animals. The control group animals were not treated. Clinical status, some rumen contents, haematological and blood biochemical parameters were examined on days 1, 3 and 7. This study has been carried out in the framework of ethics rules of Afyon Kocatepe University Ethical Committee of Animal Experiments with the reference number of AKUHA-DYEK 503-15, and was supported by Afyon Kocatepe University Scientific Research Projects Coordination Unit (BAPK) the 16.SAĞBİL.13.

Clinical evaluation

Clinical examinations such as body temperature, respiration and heart rate, and number of rumen contractions were performed according to the method reported by Hungate (26).

Hematological evaluation

Blood samples were taken from jugular venous in all animals, and hematological indices such as erythrocyte (RBC) and total leukocyte count (WBC), hematocrit (HCT), hemoglobin (HGB) level, mean corpuscular volume (MCV), mean corpuscular hemoglobin (MCH), mean corpuscular hemoglobin concentration (MCHC) were measured using commercial kits and a blood cell counter (Mindray BC2800 Vet).

Serum biochemical parameters

Serum aspartate aminotransferase (AST), alkaline phosphatase (ALP), lactate dehydrogenase (LDH), ornithine carbamoyl transferase (OCT), sorbitol dehydrogenase (SDH) were measured spectrophotometrically (SHIMADZU UV). Total proteins (TP), creatinine (CREA), albumin (ALB),

glucose (GLU), total bilirubin (TB), and direct bilirubin (DB) were measured by an autoanalyzer (Roche brand Cobas C111 Model) using commercial kits.

Rumen fluid analyzes

Methylene blue test, total infusoria count, and sedimentation test along with pH values (Mulristix 10 SG-Bayer®-Germany) were measured in fresh rumen contents taken by rumen tubes according to Boyne (27).

Statistical analyzes

Statistical calculations were made according to the ANOVA method. The Duncan test was used to determine the difference between groups. Statistical analyzes were performed using the Windowscompatible SPSS 18.0 (Inc., Chicago, II, USA) package program. Data were presented as mean \pm standard error, and P< 0.05 was considered as statistically significant.

Results

Ten out of 20 animals assigned to the case group were female and the remaining were male. The mean age of the case group was 13.2 ± 3 months. In the control group, 3 out of 10 cattles were female. The mean age of control group animals was 13.3 ± 2 months, and there was no statistically significant difference between the two groups in terms of age (P>0.05).

Clinical findings

No significant difference was found with regard to body temperature, respiration, and heart rates between the groups. However, with respect to the ruminal movements at 5 minutes, the mean values of the rumen movement in the case group were significantly higher than the control group (P< 0.05), although the values obtained were within normal limits (Table 1).

Ruminal fluid findings

The mean number of infusoria reached the highest level on the 7^{th} day ($320.30\pm104.90~\text{mm}^3$), and these higher levels were significantly higher than the infusoria counts of the control and the other working groups (P<0.05). On the contrary, in both case and control groups the methylene blue average times on the 7^{th} day were lower than the first day of the case group (Table 2).

Hematological findings

Mean WBC and RBC values of the case group were found to be significantly higher (P<0.05) when compared to those of the control group. The mean HGB levels measured on days 3 and 7 in the case group were found to be significantly higher (P < 0.05) when compared to those of the first day, and the control group (12.38 \pm 1.10 g/dL and 8.16 \pm 1.10 g/dL, respectively). The mean MCV value on 7th day in the case group was significantly higher than those of the control group, and the 1st and 3rd days. Similarly, the MCHC levels of the study group were significantly higher (P<0.05) than the control group. Furthermore, the average number of neutrophils on the control group was significantly lower (P<0.05) than the average of the case group (Table 3).

Metabolic profile findings

Table 1. Comparison of clinical features of the animals										
Groups/Parameters	Control	1st Day	3 rd Day	7 th Day	Reference values*	P				
Temperature (°C)	38,81± 1.42	38.74± 1.66	38.60± 1.67	38.56± 1.44	37.5-38.6	> 0.05				
Pulse (beats/min)	94± 6	95± 5	96± 6	96± 5	70-120	> 0.05				
Respiration (breaths/min)	26± 3	27± 2	27± 3	26± 3	15-30	> 0.05				
Rumen movement (contractions/5 min)	7± 2°	10± 2 ^b	10± 3 ^b	12± 2ª	6-12	< 0.05				

 $a,\!b,\!c \ Different \ letters \ within \ the \ same \ column \ indicate \ statistically \ significant \ differences \ between the mean \ values \ of \ the \ groups \ (P<0.05).$

The AST, ALP and SDH enzyme levels in the case group were found to be higher than the control group (P< 0.05), but the LDH and OCT levels did not differ significantly (P> 0.05). The case group urea level was found to be at the highest level $(46.32\pm6.04 \text{ mg/dL})$ on day 3, though within the

reference limits. TP, ALB, CREA, TB, and DB mean concentrations were found to be significantly higher (P < 0.05) than the control group. The GLU concentration was interestingly at the lowest level $(4.76 \pm 1.32 \text{ mg/dL})$ on day 3 (Table 4).

Groups	pH	in the control and study grou Infusoria (mm³)	Methylen Blue Test (min)	Sedimentation Test (min)
Control	7.00± 0.00	162,80+ 60,40 ^b	3.50±0.50 ^a	3.15±0.50 ^b
1st Day	7.00± 0.00	250.30± 90.20bc	3.50±0.40°	4.10±0.50°
3 rd Day	7.10± 0.00	230.60± 88.80 ^b	2.50±0.40 ^b	4.20±0.50 ^a
7 th Day	7.10± 0.00	320.30± 104.90a	2.50±0.40b	4.30±0.50 ^a
P	> 0.05	< 0.05	p<0.05	p<0.05

0.05).

Table 3.	. Hematolo	gical findin	gs and statis	tical analys	is results								
Groups	\mathbf{WBC} $(10^3/\mu\mathrm{L})$	RBC (10 ⁶ /μL)	HGB (g/dL)	HCT (%)	MCV (fl)	MCHC (g/dL)	$ \mathbf{PLT} \\ (10^3/\mu\mathrm{L}) $	MCH (pg)	Lymphocytes $(10^3/\mu\text{L})$	Monocytes $(10^3/\mu\text{L})$	Neutrophils $(10^3/\mu L)$	Eosinophils $(10^3/\mu L)$	Basophils $(10^3/\mu\text{L})$
Control	8.20± 2.20 ^b	$5.10\pm 0.60^{\pm}$	$\begin{array}{c} 8.16\pm\\ 1.10^{\mathrm{b}} \end{array}$	26.80± 5.00 ^b	52.67± 8.20°	$30.40\pm$ 2.60°	226.80 ± 19.48^{c}	16.10± 1.76 ^b	4.40± 0.40	0.72± 0.04	5.20 ± 1.80^{b}	1.48± 0.40	1
1st Day	9.80 ± 2.40^{a}	$6.10\pm 0.40^{\pm}$	$12.38\pm\\1.10^{a}$	$38.40\pm 4.60^{\pm}$	$62.94\pm\\9.60^{ab}$	$\begin{array}{c} 32.28 \pm \\ 2.42^b \end{array}$	$240.30\pm\\34.30^{b}$	$\begin{array}{c} 20.22 \pm \\ 2.64^{a} \end{array}$	4.30± 0.60	0.70 ± 0.03	6.33 ± 2.18^{a}	$1.65\pm\\0.80$	ı
3 rd Day	9.70 ± 2.60^{a}	6.30 ± 0.60^{a}	13.30 ± 2.10^{a}	$39.10\pm$ 5.40^{a}	$62.06\pm\\9.88^{ab}$	$34.20\pm$ 3.68^{a}	242.12 ±	21.10 ± 3.06^{a}	4.10± 0.80	0.70± 0.04	6.24 ± 2.30^{a}	1.63± 0.46	1
7 th Day	9.90 ± 2.30^{a}	6.20 ± 0.60^{a}	13.60 ± 1.10^{a}	39.09 ± 4.80^{a}	64.40 ± 8.90^{a}	34.68 ± 2.38^{a}	$284.28\pm$ 43.26^{a}	$\begin{array}{c} 21.78 \pm \\ 2.20^{a} \end{array}$	4.30± 0.30	$\begin{array}{c} 0.72 \pm \\ 0.02 \end{array}$	6.32 ± 2.18^{a}	1.62 ± 0.66	1
Reference Values*	6-12	5-8	8-14	24-46	37-54	26-36	175-620	11-17	3-7.5	0.1-1.5	1.5-5	0.1–1.5	×
Ь	< 0.05	< 0.05	< 0.05	< 0.05	< 0.05	< 0.05	< 0.05	< 0.05	> 0.05	> 0.05	< 0.05	> 0.05	> 0.05

a,b,c Different letters within the same column indicate statistically significant differences between the mean values of the groups (P< 0.05) R: rarely; * Filder (25).

Table 4	4. Comp	arison o	of the me	tabolic p	rofile para	ameters						
Groups	AST (III/I.)	SDH (TI/IL)	ALP (IU/L)	LDH (IU/L)	OCT (IU/L)	UREA (mg/dL)	CREA (mg/dL)	TP (g/dL)	ALB (g/dL)	GLU (mg/dL)	TB (mg/dL)	DB (mg/dL)
Contr	$116.13\pm24.40^{\circ}$ 102.28 ± 20.10^{d}	9.12±2.04b	121.32±23.18 ^b	1240.20±88.30 1246.78±90.52	178.20±12.28	40.12±50.22 ^b	$1.22\pm0.14^{\rm b}$	6.15 ± 0.10^{b}	2.94 ± 0.42^{b}	6.72 ± 1.28^{a}	0.32 ± 0.03^{d}	0.08 ± 0.001^{d}
1st Day	$116.13\pm24.40^{\circ}$	10.44 ± 2.20^{ab}	134.32±40.40ª 132.42±36.26ª	1240.20±88.30	177.30±10.10 176.68±14.24	45.24 ± 5.10^{a}	1.86 ± 0.20^{a}	6.96±1.12ªb	$3.40\pm0.42^{ m ab}$	5.45 ± 1.30^{b}	$0.56\pm0.02^{\circ}$	0.10± 0.001°
3rd Day	$123.43\pm$ 32.22^a	11.24 ± 1.60^{a}	134.32 ± 40.40^{a}	1254.16± 72.12	177.30±10.10	46.32 ± 6.04^{a}	2.02 ± 0.48^{a}	7.84 ± 1.40^{a}	4.24 ± 0.38^{a}	4.76±1.32°	0.70 ± 0.02^{a}	0.13 ± 0.001^{a}
7 th Day	121.68±36.20b 123.43± 32.22a	$11.32{\pm}1.88^a$	136.42±38.30⁴	$1260.44 \pm 80.00 1254.16 \pm 72.12$	183.22±14.20	45.24 ± 6.12^{a}	$1.96\pm0.32^{\mathrm{a}}$	$7.56 \pm 1.28^{\mathrm{a}}$	4.36 ± 0.26^a	5.12 ± 1.44^{b}	$0.62\pm0.01^{\mathrm{b}}$	0.12 ± 0.001^{b}
Reference values	78-132	4.3-15.3	90-170	<1500	205±135	42.8-64.2	1-2	6.7-7.5	3.0-3.6	4.5-7.5	0.1-0.5	0.04-0.14
Ъ	< 0.05	< 0.05	< 0.05	> 0.05	> 0.05	< 0.05	< 0.05	< 0.05	< 0.05	< 0.05	< 0.05	< 0.05

a,b,c Different letters within the same column indicate statistically significant differences between the mean values of the groups (P<0.05).

Discussion

Although ruminants are able to produce their own B complex vitamins by microbial synthesis in their rumen, some recent researchs have shown that ruminal synthesis of B complex vitamins may not be sufficient for an optimal production and health of today's cattle (especially dairy cattle) (12, 24, 25). Most B vitamins' microbial synthesis is increased by energy uptake, and decreased in the presence of B vitamins. In addition, there is a problem of complete destruction of the rumen which is added to the baits (4, 6). It has also been reported that the addition of vitamin B12 to food does not lead to a significant increase in blood B12 levels in the absence of methionine (25). A similar situation exists in cobalt (Co) deficiency (24, 28). This is why in the present study we used B complex vitamins by IM route.

Moreover, studies have shown that injections of B complex vitamins are more effective than tablets, gels or liquids (22, 29-31).

In the present study, body temperature, respiration and heart rates were not significantly different from those of the control group, but the average of the case group was higher than the control group in terms of the number of rumen contractions at 5 minutes. These findings are consistent with those reported by other researchs showing that B complex vitamins stimulate the nervous system, and might lead to muscle contractions increase, and consequently higher rumen motility (32, 33).

Furthermore, in the present study, rumen total infusoria counts on the 7th day were higher than the mean values obtained in the control group or other

time periods (days 1 and 3 after the beginning of the study), while sedimentation test averages were short. However, some researchs where B complex vitamins were administered intramuscularly led to significant positive changes on the ruminal microflora and fauna (12, 34).

The mean values of the WBC, RBC, neutrophils, HGB, HTC, MCV, MCH and MHCH in the case group were higher compared with those of the control group, but the mean values of lymphocytes, monocytes, and basophils were not different from those of the control group. Smilar findings were obtained by Uslu et al. (35) who reported that immunocapacitance increased after application of B complex vitamins.

We found that AST, SDH, and ALP enzyme levels in the case group were found to be significantly higher when compared with those of the control group, while being within the reference limits. A possible reason for this condition may be that B complex vitamins cause an increase in liver and kidney metabolism. In fact, B complex vitamins stimulate fat and protein metabolism (13). Similarly, higher urea and CREA levels can be attributed to increased ruminal fauna, and renal metabolic activity (3). It has also been reported that despite the ammonia flow in animals, nitrite conversion may decrease, resulting in an increase in blood urea levels (36). However, the increase in the concentration of CREA following the application of B complex vitamins has been adhered to increase the reabsorption activity of the kidneys (from the tubules), and possible irritation of the kidney tissue under this condition (37).

In the present study, we detected higher TP and ALB concentrations in cases compared to those of the control group. These findings were compatible with some other reports (38-40).

B vitamins, help particularly to convert foods taken by the body (vitamin B12) into glucose that can be further used by the body (24, 41, 42). In the present study, the lowest GLU concentration on day $3 (4.76 \pm 1.32 \text{ mg/dL})$ might be due to the fact that B

complex vitamins caused an increase in glucose utilization and catabolism. However, a quite large amount of B complex vitamins was needed for conversion of VFA's into energy (43).

Concentrations of TB and DB in our study were at the highest level on the 3rd day. This can be interpreted as a sign of an increase inliver metabolic activities as some researchers' reported previously (44-46). Correspondingly, some studies have shown that TB and DB concentrations can increase with the increase of liver and bile system activity (47-49).

Consequently, the use of B complex vitamins by İM route at certain intervals in cattle has proven to be of great benefit as it was not affected by the negative effects of their digestive system.

Conflict of interest

The authors declare that they have no competing interest.

References

- Coleman G S, Rumen Ciliate Protozoa, in Advances in Parasitology, W.H.R. Lumsden, R. Muller, and J.R. Baker, Editors, Academic Press. 1980:121-73.
- Demeyer D I. Rumen microbes and digestion of plant cell walls. Agriculture and Environment. 1981;6:295-337.
- Castillo-González A, Burrola-Barraza M, Domínguez-Viveros J, et al. Microorganismos y fermentación ruminal. Archivos de medicina veterinaria. 2014;46:349-61 [Article in Spanish].
- Lardinois C C, Mills R C, Elvehjem C A, et al. Rumen Synthesis of the Vitamin B Complex as Influenced by Ration Composition. J Dairy Sci. 1944;27:579-83.
- 5. Elliot J, Propionate metabolism and vitamin B12, in Digestive physiology and metabolism in ruminants. Springer. 1980:485-503.
- Leklem JE. Vitamin B6. In: Handbook of Vitamins (Machlin LJ. ed, Marcel Dekker, Inc., New York:1991;341-65.
- Girard C. B-complex vitamins for dairy cows: A new approach.
 Can J Anim Sci. 1998;78:71-90.
- 8. Elmadfa I, Majchrzak D, Rust P, et al. The thiamine status of adult humans depends on carbohydrate intake. Int J Vitam Nutr Res. 2001;71:217-21.
- Zeisel S H. Choline deficiency. J Nutr Biochem. 1990;1:332-49.

- 10. Kim Y I, Miller J W, Da Costa K A, et al. Severe folate deficiency causes secondary depletion of choline and phosphocholine in rat liver. J Nutr. 1994;124:2197-203.
- 11. Mcdowell L R, Vitamins in animal nutrition: comparative aspects to human nutrition. Elsevier: 2012.
- Santschi D. Rate of B-complex vitamins in the gastrointestmal tract of dairy cows. Doctorate Thesis. McGill University Montreal, Quebec, Canada, 2004.
- Radostits O, Gay C, Hinchcliff K, et al. A textbook of the diseases of cattle, sheep, goats, pigs and horses. Veterinary Medicine 10th edition Bailliere, Tindall, London, UK. 2007;1576-80.
- 14. Erickson P, Murphy M, Mcsweeney C, et al. Niacin Absorption from the Rumen1. J Dairy Sci. 1991;74:3492-5.
- NRC (National Research Council). Mineral Tolerance of Animals (2nd rev. ed.) National Academy of Sciences-national Research Council, Washington, D.C., USA, 2005.
- 16. Rohlfs E M, Garner S C, Mar M-H, et al. Glycerophosphocholine and phosphocholine are the major choline metabolites in rat milk. J Nutr. 1993;123:1762-8.
- 17. Virtanen A I. Milk production of cows on protein-free feed. Science. 1966;153:1603-14.
- NRC (National Research Council). Vitamin Tolerance of Animals. National Academy of Sciences-National Research Council, Washington, D.C., USA, 1987.
- NRC (National Research Council). Nutrient Requirements of Domestic Animals: Nutrient Requirements of Dairy Cattle (Sixth Rev. Ed.). National Academy of Sciences-National Research Council, Washington, D.C., USA, 1989.
- Guyton A C and Hall J E. Textbook of Medical Physiology.
 Philadelphia, Pennsylvania. 2006;1248.
- 21. Bender D A. Optimum nutrition: thiamin, biotin and pantothenate. Proc Nutr Soc. 1999;58:427-33.
- 22. Akins M S, Bertics S, Socha M, et al. Effects of cobalt supplementation and vitamin B 12 injections on lactation performance and metabolism of Holstein dairy cows. J Dairy Sci. 2013;96:1755-68.
- Fileder SE. Hematologic Referance Ranges. In: Merck Veterinary Manuel. Ailleo, SE, 11 Ed. Merck & Co., Inc., Kenilworth, NJ, USA.
- 24. Lin Y-H, Wu J-Y, Shiau S-Y. Dietary cobalt can promote gastrointestinal bacterial production of vitamin B 12 in sufficient amounts to supply growth requirements of grouper, Epinephelus

- malabaricus. Aquaculture. 2010;302:89-93.
- 25. Safarkhanlou M, Maheri-Sis N, Ebrahimnezhad Y, et al. The Effects of Intramuscular Injections of Vitamin B12 on Production Performance of Dairy Cows in Early Lactation Fed Dietary Supplements of Rumen-Protected Methionine. J Biol Environ Sci. 2016;10:11-9.
- 26. Hungate R E. The rumen and its microbes Academic Press

 New York and London 1966
- 27. Boyne A, Eadie J M, Raitt K. The development and testing of a method of counting rumen ciliate protozoa. Microbiology. 1957:17:414-23.
- Latteur J. Cobalt deficiencies and sub-deficiencies in ruminants. Cobalt deficiencies and sub-deficiencies in ruminants. 1962:1-48
- 29. Letendre M, Bernier J F, Girard C L, et al. Effects of intramuscular injections of folic acid on folates status and growth performance of weanling pigs. Can J Anim Sci. 1991;71:1223-31.
- 30. Lobley G, Connell A, Revell D. The importance of transmethylation reactions to methionine metabolism in sheep: effects of supplementation with creatine and choline. Br J Nutr. 1996;75:47-56.
- 31. Luce W, Buchanan D, Maxwell C, et al. Effect of supplemental choline and dichlorvos on reproductive performance of gilts. Nutr Rep Int. 1985;32:245-51.
- 32. Garosi L S, Dennis R, Platt S R, et al. Thiamine deficiency in a dog: clinical, clinicopathologic, and magnetic resonance imaging findings. J Vet Intern Med. 2003;17:719-23.
- Gibson G E and Zhang H. Interactions of oxidative stress with thiamine homeostasis promote neurodegeneration. Neurochem Int. 2002;40:493-504
- 34. Cole N, Mclaren J, Hutcheson D. Influence of preweaning and B-vitamin supplementation of the feedlot receiving diet on calves subjected to marketing and transit stress. J Anim Sci. 1982:54:911-7.
- 35. Uslu U, Balevİ T, UÇan U S, et al. Effect of parenteral administration of vitamin B to goats on performance, lice (Phthiraptera) infestations and cellular immunity. Kafkas Üniversitesi Veteriner Fakültesi Dergisi. 2017;23:715-20.
- 36. Valente T N P, Da Silva Lima E, Dos Santos W B R, et al. Ruminal microorganism consideration and protein used in the metabolism of the ruminants: A Review. Afr. J. Microbiol. Res. 2016;10:456-64.

Effects of Vit B Complex on Ruminal Protozoa

- 37. Taes Y E, Delanghe J R, De Vriese a S, et al. Creatine supplementation decreases homocysteine in an animal model of uremia. Kidney Int. 2003;64:1331-7.
- 38. Jouany J and Ushida K. The role of protozoa in feed digestion-Review. Asian-Australas J Anim Sci. 1999;12:113-28.
- Gräsbeck R. Imerslund-Gräsbeck syndrome (selective vitamin B 12 malabsorption with proteinuria). Orphanet J Rare Dis. 2006;1:17.
- Storm T, Emma F, Verroust P J, et al. A patient with cubilin deficiency. N Engl J Med. 2011;364:89-91.
- 41. Latimer K S, Duncan and Prasse's veterinary laboratory medicine: clinical pathology. John Wiley & Sons:2011.
- 42. Zinn R, Owens F, Stuart R, et al. B-vitamin supplementation of diets for feedlot calves. J Anim Sci. 1987;65:267-77.
- 43. Mcsherry B, Lumsden J, Valli V, et al. Hyperbilirubinemia in sick cattle. Can J Comp Med. 1984;48:237.
- 44. Smith BP. Large Animal Internal Medicine. Mosby Elsevier,

- St. Louis, MO, USA, 2009.
- 45. Allison R. Laboratory evaluation of the liver. Veterinary Hematology and Clinical Chemistry. 2nd ed. Ames, IA, USA: Wiley-Blackwell Inc. 2012;401-24.
- 46. Dubeski P, Owens F, Song W, et al. Effects of B vitamin injections on plasma B vitamin concentrations of feed-restricted beef calves infected with bovine herpesvirus-1. J Anim Sci. 1996:74:1358-66.
- 47. Elİtok B, Elİtok Ö M, AkgÜn S. B Vitaminleri Yetmezliği. Turkiye Klinikleri Journal of Veterinary Sciences-Internal Medicine-Special Topics. 2016;2:51-5 [Article in Turkish].
- 48. Harmeyer J and Kollenkirchen U. Thiamin and niacin in ruminant nutrition. Nutr Res Rev. 1989;2:201-25.
- 49. Silzell S, Hellwig D, Kegley E, et al. Effects of supplemental thiamin on growth performance and immune function in stressed stocker cattle. J Appl Anim Res. 2002;22:145-56.